arXiv:1411.2353v1 [nucl-th] 10 Nov 2014

^{136}Sn and three body forces

M. Saha Sarkar [*], **S. Sarkar** [†]
Saha Institute of Nuclear Physics, Bidhannagar, Kolkata-700064, INDIA
Indian Institute of Engineering Science and Technology, Shibpur, Howrah - 711103, INDIA

November 11, 2014

Abstract

New experimental data on 2^+ energies of $^{136,138}Sn$ confirms the trend of lower 2^+ excitation energies of even-even tin isotopes with N $>$ 82 compared to those with N$<$ 82. However, none of the theoretical predictions using both realistic and empirical interactions can reproduce experimental data on excitation energies as well as the transition probabilities ($B(E2; 6^+ \rightarrow 4^+)$) of these nuclei, simultaneously, apart from one whose matrix elements have been changed empirically to produce mixed seniority states by weakening pairing. We have shown that the experimental result also shows good agreement with the theory in which three body forces have been included in a realistic interaction. The new theoretical results on transition probabilities have been discussed to identify the experimental quantities which will clearly distinguish between different views.

1 Introduction

Nuclei around doubly closed ^{132}Sn lie on or close to the path of astrophysical r-process flow. Structure of these nuclei, particularly the binding energy (BE), low-lying excited states and beta decay rates at finite temperatures are important ingredients for nucleosynthesis calculation. However, these nuclei are usually experimentally inaccessible by common techniques of nuclear spectroscopy. So far experimental investigations have been performed using spontaneous fission sources, thermal - neutron - induced fission deep inelastic reactions, fragmentation and fission at intermediate and relativistic energies. Reactions with neutron rich radioactive ion beams are expected to generate very important, reliable data base for these nuclei.

So far the experimental status [1] was not at all satisfactory for Sn isotopes beyond ^{132}Sn. Spectroscopic information, such as BE and low lying spectrum, is known experimentally only for ^{134}Sn. Half-lives of $^{135-137}Sn$ have been measured through β-n decay process. Lifetimes of these nuclei are very small and production rates are also very low presenting challenges to spectroscopic studies. Reliable theoretical results are therefore necessary and useful, especially as inputs to calculations for astrophysical processes. Being close to the doubly closed neutron-rich ^{132}Sn nucleus, this region therefore has been taken as a fertile ground for testing nuclear shell model (SM) with interactions suited for nuclei far from stability.

[*]maitrayee.sahasarkar@saha.ac.in
[†]ss@physics.becs.ac.in

THE CORE AND THE VALENCE SPACE



Figure 1: The valence space for proton (p) and neutrons (n) above ^{132}Sn core.

2 Evolution of our endeavours

We have started working in this mass region in early 2000, with a systematic study [2] of this mass region utilising available interactions. We have used then available version of the shell model code OXBASH [3]. The interactions [4] used by us were KH5082 and CW5082. The SM calculations were carried out in the Z50N82 model space (Fig. 1). The earlier workers realised that there are many points of similarity between the spectroscopy of the doubly closed shell regions around stable ^{208}Pb and neutron-rich ^{132}Sn. The two interactions mentioned above are such (1+2)- body nuclear interaction Hamiltonians where interactions of ^{208}Pb region are scaled down to the next lower available doubly magic domain, i.e., in the ^{132}Sn region. Most of our results satisfactorily explained the available experimental data for several neutron rich isotopes of $_{52}Te$, $_{53}I$, $_{54}Xe$, $_{55}Cs$, $_{56}Ba$ with N=82 and 83. However, we found for N$\geq$84 isotones, the interactions were not as successful and seemed to be inappropriate. The evergy levels were overpredicted for these neutron rich nuclei (Fig. 2). So the limitations of the two interactions in predicting experimental observables in N$\geq$84 isotones, clearly indicated a necessity of changing particularly the $n-n$ two body matrix elements (tbmes) of the interactions.

2.1 Construction of the new Hamiltonian

Keeping the conclusion of our first work in this mass region in mind, we attempted a simple modification of the CW5082 interaction [4]. The CW5082 Hamiltonian was modified in the light of available information on binding energies, low-lying spectra of two-nucleon nuclei: A=134 isobars of Sn,Sb and Te isotopes. The energies of the single particle orbitals of the valence space above the ^{132}Sn core have been replaced by the experimental ones. The details of this modification procedure have been discussed in Ref. [5]. The new Hamiltonians work remarkably well in predicting binding energies, low-lying spectra and electromagnetic transition probabilities for N=82,83 and even for N $\geq$ 84 isotones of Sn,Sb,Te,I,Xe and Cs nuclei. Out of total two thousand one hundred and one (2101) tbmes, only twenty six (26) tbmes are modified in SMPN interaction. Twelve (12) of them are $n-p$ tbmes, four (4) are $p-p$ and the rest are ten (10) are $n-n$ tbmes. Six of these neutron-neutron matrix elements are diagonal elements coupled to 0^+ spin. These matrix elements are reduced, implying a reduction of the pairing this in neutron-rich domain. The modified two body matrix elements of SMPN are tabulated in Ref.[6].

2.2 Comparison with realistic interaction

The success of his new interaction prompted us to apply it for several new data sets obtained by experimentalists [7, 8, 9]. In the mean time, few other groups have worked with CD-Bonn potential based realistic interctions [10, 11, 12]. Their results also agreed well with many of the available data. So our next aim was to compare the results obtained with the empirically modified inteactions with those obtained with the realistic interactions.

We have used the CD-Bonn potential based realistic CWG [10] and the empirical SMPN interactions. These Hamiltonians are used in this neutron-rich region and have the same set of single-particle energies of the valence orbitals but different sets of two-body interaction matrix elements. The SM calculations [13] were carried out in the Z50N82 model space (Fig. 1) with the SMPN and CWG Hamiltonians using the code OXBASH and NuShell [3].

The calculations have been done [13] for ^{138}Sn (T_z=+3.0) with 6 neutrons, ^{138}Te (T_z=+1.0) with 4 neutrons and 2 protons, ^{138}Xe (T_z=-1.0) with 2 neutrons and 4 protons and finally, ^{138}Ba (T_z=-3.0) with 6 valence protons. Untruncated shell model calculations for 6 valence nucleons in



Figure 2: Comparison of experimental and theoretical energy spectra of ^{136}Te.

this basis space usually involve matrices of large dimensions. Calculated energy eigenvalues of the levels with the SMPN and CWG Hamiltonians are in good agreement with experimental ones up to the highest observed level in ^{138}Te (Fig. 3), ^{138}Xe and ^{138}Ba [13] (including the non-yrast levels in $^{138}Xe, Ba$). In the CWG results all the energy eigenvalues of the excited states are systematically underpredicted by about 100-150 keV. For neutron rich ones like ^{138}Te, both these calculations reproduced quite reasonably well the vibrational spectrum of this nuclei (Fig. 3) indicated by its R_4= E(4_1^+)/E(2_1^+) = 2.04 value. In both the results, the lowest proton and neutron orbitals have the maximum occupations. The CWG wavefunctions have comparatively larger occupations in other orbitals also, indicating larger configuration mixing.

2.3 Predictions for neutron rich heavy Sn isotopes

It was remarkable that for $^{136,138}Sn$, where the experimental level schemes were not known until recently, the two results [13] differ dramatically (Figs. 4,6). The energies of the first three excited states 2_1^+, 4_1^+ and 6_1^+, for ^{138}Sn with the CWG Hamiltonian appear at about twice of those predicted by SMPN.

In ^{138}Sn, multiplet structure with $\nu 2f_{7/2}^6$ is more clear in SMPN results for 0^+ to 6^+ states. Six neutrons or in other words two neutron holes in $\nu 2f_{7/2}$ can couple to $J_{max} = 6$. Thus for generating the 8^+ state a pair of neutrons must be promoted to a higher lying single particle orbital. This leads to an energy gap of about 2.0 MeV, in the spectrum between the first 6^+ and 8^+ states, as shown in the SMPN results (Fig.6). This gap is about 0.8 MeV for CWG.

While comparing [13] the structure of the states in ^{138}Ba with six valence protons with that of the ^{138}Sn with the same number of neutrons, it is found that the protons in this valence space are more efficient in generating configuration mixing. This is essentially due to close spacing between



Figure 3: Comparison of experimental data and theoretical predictions for excitation energies in ^{138}Te.



Figure 4: The calculated excitation spectra of ^{136}Sn using SMPN, CWG3M and CWG interactions. Results have been compared with experimental data.

$\pi(gds)$ single particle orbitals that permits $\pi - \pi$ residual interaction to scatter easily protons to various single particle states.

The wave function structure shows [13, 14] that the CWG interaction favors large configuration mixing conserving seniority as far as possible. In contrast, SMPN favors the purer structure of the low-lying states and shows characteristics of $\nu(2f_{7/2})$ multiplets. Interestingly, this structural difference has a dramatic effect in the transition rates (especially, $B(E2; 2^+ \rightarrow 0^+)$ calculated with these two interactions for both $^{136,138}Sn$ [14].

In Ref. [13], energies,wave functions, and $B(E2, 0^+_{g.s.} \rightarrow 2^+_1)$ values have been compared for the yrast 0^+ and 2^+ states of the three Sn isotopes. It was found that although the $E(2^+_1)$ energies decrease with increasing neutron number in Sn isotopes for N$\geq$84, B(E2) values corresponding to the SMPN calculations do not increase as expected. This feature was already observed experimentally in the neutron-rich ^{136}Te isotope in this mass region. Origin of an anomalously low $B(E2, 0^+_{g.s.} \rightarrow 2^+_1)$ value found in this nucleus was to traced to a reduced neutron pairing above the N = 82 shell gap. Interestingly, similar behavior is also seen near the N = 20 shell closure for neutron-rich Mg and Ne isotopes [13]. It seems to be a unique feature observed in neutron-rich nuclei. In this domain, a comparatively lower value of $E(2^+_1)$ does not necessarily lead to a larger B(E2) value indicating collectivity. This type of feature is predicted theoretically with the SMPN interaction for N $\geq$ 84.

2.3.1 The shell closure at N=90 for Sn isotope and its origin

In our next work [14], with SMPN interaction, the prediction for E(2_1^+) of ^{140}Sn comes out to be much larger than the expectation. The comparison of the high 2_1^+ energy of ^{140}Sn with examples from other neutron-rich domains [14] clearly show that N = 84-88 spectra with SMPN manifest the effect of gradually filling up the $\nu(2f_{7/2})$ orbital, which finally culminates in a new shell closure at N = 90.

In order to understand how the shell closure at N=90 develops as one increases number of valence neutrons from ^{132}Sn, the effective single-particle energies (ESPE) for the neutron orbitals for the two Hamiltonians have been compared [14]. To investigate the specific component of the interaction which is responsible to generate this new shell gap, the two body matrix elements of both CWG and SMPN have been spin-tensor decomposed [14, 15] in to central, antisymmetric spin- orbit (ALS), spin-orbit (LS) and tensor parts [14]. For SMPN, the central and ALS part for $2f_{7/2}$- $2f_{7/2}$ tbmes account for majority of the downward shift of the ESPE of $2f_{7/2}$ with increasing valence neutron number (n). Variation in ALS part is primarily responsible for this observed shell gap at N=90 [14].

While analysing our results, we have to understand the implication of ALS term. Bare nucleon-nucleon force contains no ALS term. A characteristic feature common to many empirical effective interactions is the strong ALS components in the tbmes. It indicates important contributions from higher order renormalisation or many body effects to the effective interactions. In empirical SMPN such many - body effects might have been included in some way through the modification of important tbmes.

However, it has been observed experimentally that N = 90 is suitable for onset of deformation for nuclei above Sn (like Xe, Ba, etc.). Therefore, the energy of the deformed configuration also comes down. The presence of valence protons above the inert core is essential for onset of collectivity. The ESPEs for SMPN which indicate the features of a shell closure for Z = 50 at N = 90, have also been analysed to indicate the possibility of onset of deformation at N = 90 with increasing Z. If the ESPEs for proton orbitals for N = 90 are plotted, substantial reduction of the $1g_{7/2}$ and $2d_{5/2}$ energy gap is observed with increasing Z, which may favor onset of collectivity [14].

3 The three body forces and its implication

It has been discussed in our earlier work ([14, 15], and references therein) that the realistic interactions fail to reproduce some shell closures for neutron-rich nuclei. It has been shown [16] that a three-body delta-hole mechanism can explain these shell gaps and three-body forces are necessary to explain why the doubly-magic ^{24}O nucleus is the heaviest oxygen isotope.

Zuker [17] had demonstrated earlier that a very simple three-body monopole term can eliminate the limitations of the realistic two-body potentials. According his prescription, we have included corrections in the relevant tbmes of CWG to incorporate the three body effects. The correction factor will be effective for nuclei for which the valence neutron number n= 3 or more. It is amazing that although CWG does not predict a shell closure at ^{140}Sn, the updated CWG including three-body forces, named as CWG3M, also predicts a shell gap for N=90. The E(2_1^+) energy of ^{140}Sn predicted by CWG3M (1.889 MeV) is close to that predicted by SMPN (1.949MeV) [14, 15, 18]. This also indicates that three body effect plays an important role for shell evolution in neutron rich Sn isotopes above ^{132}Sn, as also observed in *sd* and *pf* shells.



Figure 5: The calculated 2^+ energies of $^{134-140}Sn$ using SMPN, CWG3M and CWG interactions. Results have been compared with experimental data.

4 The new results on $^{136,138}Sn$ and their interpretations

Very recently, experiments were carried out at the RIKEN Radioactive Isotope Beam Factory (RIBF) [19, 20] to study the neutron rich isotopes of Sn. In one of the experiments the first 2^+ excited state in the neutron-rich tin isotope ^{136}Sn has been identified at 682(13) keV by measuring γ -rays in coincidence with the one proton removal channel from ^{137}Sb [19]. In another experiment, delayed γ-ray cascades, originating from the decay of isomeric states at I=6^+, in the very neutron-rich, semimagic isotopes $^{136,138}Sn$ have been observed following the projectile fission of a ^{238}U beam at RIBF, RIKEN. The measured $E_x(2_1^+)$ value of ^{136}Sn was found to be much higher than the neighboring N = 86 isotones, indicating a good Z = 50 shell closure. The excitation energies stay almost constant in ^{134}Sn, ^{136}Sn and ^{138}Sn, which suggests that the seniority - 2 coupling scheme holds beyond N = 82 up to N = 88[19]. They found that the predicted and experimental $B(E2; 6^+ \rightarrow 4^+)$ values [20] agree for ^{138}Sn with both empirical and realistic interactions. However, for ^{136}Sn the results with their realistic interaction, differ by a factor of > 5. Three other shell-model calculations, using realistic and empirical interactions, also failed to reproduce this value for ^{136}Sn by at least a factor of 2 [20]. To interpret this data better, the authors incorporated an empirical modification to the $2f_{7/2}^2$ matrix elements, probably the zero coupled ones, equivalent to reduced pairing, which generates a seniority-mixed 4_1^+ state, reproducing all available experimental data better. The authors proposed that the wave functions of these isomeric states [20] to be predominantly a fully aligned pair of $f_{7/2}$ neutrons.



Figure 6: The calculated excitation spectra of ^{138}Sn using SMPN, CWG3M and CWG interactions. Results have been compared with experimental data.

5 Motivation of the present work

From the experimental data on odd-even staggering of masses, it has been shown [21] that variation of pairing as a function of neutron number plays an important role in many distinctive features like occurrence of new shell closures, shell erosion, anomalous reduction of the energy of the first 2^+ state and slower increase of $B(E2; 2_1^+ \rightarrow 0_1^+)$ in the neutron-rich even-even nuclei of different mass regions. Recently, it has been indicated [22] that it is essential to include 3N contributions to the pairing interaction for a quantitative description of nuclear pairing gaps. It is usually found that inclusion of a three body effect improves the predictive power of realistic interactions. In our earlier work, we have already included three body effects in realistic CWG interaction to construct CWG3M. Our aim is to study the importance of three body forces in interpretation of the new data.

6 Shell Model Calculations

In the present work we have used the CD-Bonn potential based realistic CWG [10] and the empirical SMPN [5] along with CWG3M [14] interaction. All these Hamiltonians have the same set of single-particle energies of the valence orbitals but different sets of two-body interaction matrix elements. The SM calculations [13] were carried out in the Z50N82 model space using the code OXBASH [3].

I_i	I_f	CWG [10]	CWG3M [14]	SMPN[5]
^{136}Sn				
2	0	125	88	90
4	2	86	81	43
6	4	14	45	14
8	6	53	48	46
10	8	0.0097	0.0948	0.0954
12	10	91	77	69
6	4(2)	52	16	45
^{138}Sn				
2	0	184	73	69
4	2	0.3286	45	44
6	4	12	21	20
8	6	1.05	0.20	0.73
10	8	138	104	81
12	10	60	56	44

Table 1: Comparison of calculated B(E2) (e^2fm^4) values for major transitions connecting the positive parity yrast levels till 12^+ for $^{136,138}Sn$. The experimental values of $B(E2; 6_1^+ \rightarrow 4_1^+)$ for $^{136,138}Sn$ is 24(4) and $\simeq 17\ e^2fm^4$, respectively [20]. Effective neutron charge is $e_n = 0.64$

7 Results

We have compared these new results with our theoretical ones calculated using CWG, SMPN and CWG3M interactions as shown in Figs. (5,4,6). The calculated B(E2) values for all the E2 transitions connecting the yrast band till 12^+ state have also been tabulated in Table 1. To complete the information - the B(E2) for $6_1^+ \rightarrow 4_2^+$ for ^{136}Sn nucleus has also been tabulated.

7.1 Energy

If we keep in mind the usual errors obtained in theoretical shell model predictions, for ^{136}Sn, one can not choose between these three interactions. However, the predicted and experimental energy spectra for both $^{136,138}Sn$(Figs. 5,4,6) show that best agreement has been obtained with the theory in which three body forces have been included in a realistic interaction (CWG3M).

7.2 Transition probabilities

The measurement of transition probabilities and its comparison with theory, is the best way to validate a theoretical prediction. As discussed in Ref. [13, 21], the depresssed $E(2_1^+)$ energies in neutron-rich nuclei arises due to weakening of pairing, unlike near stability, where low $E(2_1^+)$ indicate collective deformed state manifested through strong B(E2). Therefore, anomalous reduction of the energy of the first 2_1^+ state and slower increase in the $B(E2; 2_1^+ \rightarrow 0_1^+)$ in the neutron-rich even-even nuclei is a distinctive feature arising out of reduced pairing. It has been discussed earlier [13], apart from differences between the $E(2_1^+)$, the most dramatic difference lies in the transition probabilities of the $2_1^+ \rightarrow 0_1^+$ transition using CWG and SMPN interactions. In the present work, it is found that the although the B(E2) values for the 6^+ isomers in $^{136,138}Sn$ are almost similar with CWG and SMPN, that from CWG3M is quite different. On the other hand, the B(E2)s for 2_1^+ states are similar for both the isotopes are similar with CWG3M and SMPN, whereas it is almost double with CWG.

8 Conclusion

The SMPN and CWG3M include the effects of reduced pairing and three body effects. It has been indicated [19, 20] that reduced pairing is responsible for the anomalous results in these nuclei. Our results clearly show that the most sensitive probe to confirm this claim will be the $B(E2; 2_1^+ \rightarrow 0_1^+)$ values. Thus the importance of pairing will be confirmed from the transition probability measurement in these neutron rich isotopes of Sn.

9 Acknowledgments

Special thanks are owed to B. A. Brown for his help in providing us the OXBASH (Windows Version) and the NUSHELL@MSU Codes.

References

[1] http://www.nndc.bnl.gov

[2] Sukhendusekhar Sarkar, M. Saha Sarkar, Phys. Rev. C **64**, 014312 (2001).

[3] B. A. Brown, A. Etchegoyen, N. S. Godwin, W. D. M. Rae, W. A. Richter, W. E. Ormand, E. K. Warburton, J. S. Winfield, L. Zhao and C. H. Zimmerman, Oxbash for Windows PC, MSU-NSCL Report No. **1289**, (2004); B. A. Brown and W. D. M. Rae,NUSHELL@MSU, MSU-NSCL Report,2007.

[4] W.T. Chou and E.K. Warburton, Phys. Rev. C **45**, 1720 (1992)

[5] Sukhendusekhar Sarkar, M. Saha Sarkar, Eur. Phys. Jour. A **21**, 61 (2004).

[6] S. Sarkar, M. Saha Sarkar, Proc. DAE-BRNS Symp. Nucl. Phys. (India) **55**, I27 (2010); (http://www.sympnp.org/proceedings/index.php: electronic version only).

[7] W. Urban, M. Saha Sarkar, S. Sarkar, T. Rzaca-Urban, J.L. Durell, A.G. Smith, J.A. Genevey, J.A. Pinston, G.S. Simpson, I. Ahmad, Euro. Phys. Jour. A **27**, 257 (2006).

[8] T. Rzaca-Urban, K. Pagowska, W. Urban, A. Zlomaniec, J. Genevey, J. A. Pinston, G. S. Simpson, M. Saha Sarkar, S. Sarkar, H. Faust, A. Scherillo, I. Tsekhanovich, R. Orlandi, J. L. Durell, A. G. Smith, and I. Ahmand, Phys. Rev. C **75**, 054319 (2007).

[9] T. Rzaca-Urban, W. Urban, M. Saha Sarkar, S. Sarkar, J.L. Durell, A.G. Smith, B.J. Varley, and I. Ahmad, Euro. Phys. Jour. A **32**, 5 (2007).

[10] B. A. Brown *et al.*, Phys. Rev. C **71**, 044317 (2005).

[11] A. Covello, L. Coraggio, A. Gargano, and N. Itaco, J. Phys.Conf. Ser. **267**, 012019 (2011).

[12] M. P. Kartamyshev, T. Engeland, M. Hjorth-Jensen, and E. Osnes, Phys. Rev. C **76**, 024313 (2007).

[13] S. Sarkar and M. Saha Sarkar, Phys. Rev. C **78**, 024308 (2008).

[14] S. Sarkar and M. Saha Sarkar, Phys. Rev. C **81**, 064328 (2010).

[15] S. Sarkar and M. Saha Sarkar, J. Phys.Conf. Ser. **267**, 012040 (2011).

[16] Takaharu Otsuka, Toshio Suzuki, Jason D. Holt, Achim Schwenk and Yoshinori Akaishi, Phys. Rev. Lett. **105**, 032501 (2010).

[17] A. P. Zuker, Phys. Rev. Lett. **90**, 042502 (2003).

[18] M. Saha Sarkar, S. Sarkar, AIP Conference Proceedings, Volume **1444**, 117 (2012).

[19] He Wang *et al.*, Prog. Theor. Exp. Phys. 023D02 (2014).

[20] G. S. Simpson *et al.*, Phys. Rev. Lett. **113**, 132502 (2014).

[21] Maitreyee Saha Sarkar, Sukhendusekhar Sarkar, Proc. of the Fifth International Conference: Sanibel Island, USA 4 -10 November 2012, Editors: J H Hamilton, A V Ramayya, World Scientific, 498 (2013).

[22] J.D. Holt, J. Menendez and A Schwenk, J. Phys. G: Nucl. Part. Phys. **40**, 075105 (2013).